



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2017

Visualization of extensive intraplaque neovascularization by optical coherence tomography

Koganti, S ; Karanasos, A ; Tu, S ; Rakhit, R D ; Regar, E

DOI: <https://doi.org/10.1016/j.hjc.2017.01.011>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-135720>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Koganti, S; Karanasos, A; Tu, S; Rakhit, R D; Regar, E (2017). Visualization of extensive intraplaque neovascularization by optical coherence tomography. *Hellenic Journal of Cardiology*, 58(1):87-88.

DOI: <https://doi.org/10.1016/j.hjc.2017.01.011>

Accepted Manuscript

Visualization of extensive intraplaque neovascularization by optical coherence tomography

S. Koganti, A. Karanasos, S. Tu, R.D. Rakhit, E. Regar

PII: S1109-9666(17)30028-3

DOI: [10.1016/j.hjc.2017.01.011](https://doi.org/10.1016/j.hjc.2017.01.011)

Reference: HJC 120

To appear in: *Hellenic Journal of Cardiology*

Received Date: 12 November 2015

Accepted Date: 2 September 2016

Please cite this article as: Koganti S, Karanasos A, Tu S, Rakhit RD, Regar E, Visualization of extensive intraplaque neovascularization by optical coherence tomography, *Hellenic Journal of Cardiology* (2017), doi: 10.1016/j.hjc.2017.01.011.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Letter to the Editor:

Title: Visualization of extensive intraplaque neovascularization by optical coherence tomography.

Koganti.S¹, Karanasos. A², Tu, S³, Rakhit R.D¹, Regar. E²

¹ Royal Free Hospital & UCL Institute of Cardiovascular Science, London

² Thoraxcentre, Erasmus MC, Rotterdam

3. School of Biomedical Engineering, Shanghai Jiao Tong University, Shanghai, China

Disclosures: None

Inflammation plays a key role in pathogenesis and progression of atherosclerosis (1). Advent of novel intravascular imaging tools such as Optical Coherence Tomography (OCT) has led to visualization of coronary atherosclerotic plaque characteristics in vivo that were previously seen only on histology. Herein we discuss a case of extensive neovascularization, a feature that imparts vulnerability to atherosclerotic plaque in the coronary artery, as seen by two- and three-dimensional OCT images.

A 63-year-old female with history of diabetes mellitus and hypertension presented with troponin-positive chest pain. She underwent coronary angiography followed by drug-eluting stent implantation to the mid segment of right coronary artery (RCA). OCT was subsequently performed to assess the post-procedural result, showing a well-deployed stent and extensive network of microvessels in a thickened intima in the proximal segment of the RCA (Figure 1). Frame-by-frame 3D reconstruction of the microvessels revealed an extensive frond-like network in the proximal segment of RCA (Figure 2).

In most patients, acute coronary syndrome (ACS) is secondary to atherosclerotic plaque rupture (2) and resultant intracoronary thrombosis. Plaque expansion secondary to intra-plaque hemorrhage (IPH) is one of the factors that can destabilize the plaque (3). IPH is thought to occur as a consequence of rupture of intra-plaque microvessels developed due to a process called neo-angiogenesis.

Human coronary artery is a tri-laminar structure and is lined by intima, media and adventitia from inside to the outside (Figure 1A&1B). The normal human coronary artery derives blood supply from vasa-vasorum situated in the adventitious layer. A process called plaque neovascularization – which involves formation of neo-vessels (Figure 1C,

1D) takes place simultaneously, initially with intimal layer thickening and later during plaque expansion. These neo-vessels originating from the vasa-vasorum reach the plaque situated in the intima through the medial layer. It has been shown that these microvessels lack structural integrity and thus are fragile and might easily rupture. Following rupture of microvessels, deposition of free cholesterol, macrophage infiltration and accumulation of erythrocyte membranes result in the enlargement of the necrotic core within an existing atherosclerotic plaque. These factors increase the risk of plaque destabilization culminating in plaque rupture and acute event.

Owing to the advent of novel intravascular imaging tools such as OCT(4,5), high-resolution pictures of atherosclerotic plaque characteristics that were once assessed ex-vivo only can now be appreciated in vivo.

References:

1. Benetos G, Toutouzas K, Drakopoulou M et al. Bilateral symmetry of local inflammatory activation in human carotid atherosclerotic plaques. *Hellenic J Cardiol*. 2015 Mar-Apr; 56(2): 118-24.
2. Virmani R, Burke AP, Farb A, Kolodgie FD. Pathology of the vulnerable plaque. *J Am Coll Cardiol*. 2006 Apr 18; 47(8 Suppl): C13-8.
3. Virmani R, Kolodgie FD, Burke AP, et al. Atherosclerotic plaque progression and vulnerability to rupture angiogenesis as a source of intraplaque hemorrhage. *Arterioscler Thromb Vasc Biol* 2005; 25: 2054–61.
4. Karanasos A, Muramatsu T, Diletti R, et al. Early and late optical coherence tomography findings following everolimus-eluting bioresorbable vascular scaffold implantation in myocardial infarction: a preliminary report. *Hellenic J Cardiol*. 2015 Mar-Apr;56(2):125-35.
5. Karanasos A, Ligthart J, Witberg K, et al. Optical Coherence Tomography: Potential Clinical Applications. *Curr Cardiovasc Imaging Rep*. 2012 Aug;5(4):206-220.




Figure 1:

1A: Normal OCT cross-section of a coronary artery

1B: Magnified view demonstrating the three layers of normal coronary artery; I – Intima, M – Media, A – Adventitia

1C: Cross-section showing microvessels

1D: Magnified view showing microvessels




Figure2:
3D reconstruction showing microvessels
from (in Red) A) Inside through cut open
view, B) Outside and C) with out coronary
artery.